



Lilly accelerating baricitinib's availability in India following receipt of permission for restricted emergency use as a COVID-19 therapy via donations and licensing agreements

May 4, 2021

- Lilly is working with the Indian government to provide baricitinib donations in conjunction with the permission
- Licensing collaboration builds upon Lilly's access principles and potentially speeds up the manufacturing and distribution of baricitinib to hospitalized COVID-19 patients requiring supplemental oxygen in India

INDIANAPOLIS, May 4, 2021 /PRNewswire/ -- Eli Lilly and Company (NYSE: LLY) announces new initiatives to help COVID-19 patients in India as part of its commitment to bring the full force of its scientific and medical expertise to attack the coronavirus pandemic around the world. Lilly is offering donations of baricitinib to the Indian government through Direct Relief while simultaneously working with local Indian pharmaceutical companies to execute royalty-free voluntary licensing agreements to accelerate the manufacturing and distribution of the medicine in India during the pandemic. An initial donation of 400,000 baricitinib tablets is being made immediately available to the Indian government for eligible hospitalized COVID-19 patients in India and Lilly will work urgently to increase the quantity of donated product multifold over the coming weeks.

On Monday, Lilly received permission for restricted emergency use from Central Drugs Standard Control Organization, a division of Ministry of Health, for baricitinib (2 mg and 4 mg) in combination with remdesivir, for treatment of suspected or laboratory confirmed coronavirus disease 2019 (COVID-19) in hospitalized adults requiring supplemental oxygen, invasive mechanical ventilation, or extracorporeal membrane oxygenation (ECMO). Baricitinib is an oral medication currently registered in India for the treatment of moderate to severe active rheumatoid arthritis in adult patients who have responded inadequately to, or who are intolerant to one or more disease-modifying anti-rheumatic drugs.

"With the COVID-19 crisis devastating India, hospitals are overwhelmed by the number of cases and patients need access to potentially life-saving treatments such as baricitinib," said David A. Ricks, Lilly chairman and CEO. "We hope that our donations as well as collaborations with other organizations speed access to baricitinib and provide treatment options for these patients."

As the global pandemic evolves, Lilly continues to evaluate opportunities to provide treatments to COVID-19 patients in countries around the world.

Authorized Use Under the EUA and Important Safety Information for baricitinib (in the United States) for COVID-19

Baricitinib is authorized for use under an Emergency Use Authorization (EUA) in combination with remdesivir, for treatment of suspected or laboratory confirmed coronavirus disease 2019 (COVID-19) in hospitalized adults and pediatric patients 2 years of age or older, requiring supplemental oxygen, invasive mechanical ventilation, or extracorporeal membrane oxygenation (ECMO).

Baricitinib has not been approved for the treatment of COVID-19, but has been authorized for emergency use by the FDA. Baricitinib is authorized under an EUA only for the duration of the declaration that circumstances exist justifying the authorization of the EUA of baricitinib under Section 564(b)(1) of the Act, 21 U.S.C. § 360bbb-3(b)(1), unless the authorization is terminated or revoked sooner.

For information on the authorized use of baricitinib and mandatory requirements under the EUA, please review the [FDA Letter of Authorization, Fact Sheet for Healthcare Providers](#) and Fact Sheet for Patients, Parents and Caregivers ([English](#); [Spanish](#)).

Important Safety Information about baricitinib for COVID-19

The following provides essential safety information on the unapproved use of baricitinib under the Emergency Use Authorization.

Warnings

Serious Infections: Serious infections have occurred in patients receiving baricitinib. Avoid the use of baricitinib with known active tuberculosis. Consider if the potential benefits outweigh the potential risks of baricitinib treatment in patients with active serious infections other than COVID-19 or chronic/recurrent infections.

Thrombosis: In hospitalized patients with COVID-19, prophylaxis for venous thromboembolism is recommended unless contraindicated. If clinical features of deep vein thrombosis or pulmonary embolism occur, patients should be evaluated promptly and treated appropriately.

Abnormal Laboratory Values: Evaluate at baseline and thereafter according to local patient management practice. Monitor closely when treating patients with abnormal baseline and post-baseline laboratory values. Follow dose adjustments as recommended in the Fact Sheet for Healthcare Providers for patients with abnormal renal, hematological and hepatic laboratory values. Manage patients according to routine clinical guidelines.

Hypersensitivity: If a serious hypersensitivity occurs, discontinue baricitinib while evaluating the potential causes of the reaction.

See **Warnings and Precautions** in the FDA-approved full Prescribing Information for additional information on risks associated with longer-term treatment with baricitinib.

Serious Side Effects: Serious venous thrombosis, including pulmonary embolism, and serious infections have been observed in COVID-19 patients treated with baricitinib and are known adverse drug reactions of baricitinib.

There are limited clinical data available for baricitinib use in coronavirus 2019 (COVID-19). Additional information regarding baricitinib for its FDA-approved indication, including safety information, may be found in the full [Prescribing Information](#), including Boxed Warning about Serious Infections, Malignancies, and Thrombosis, and [Medication Guide](#).

Use in Specific Populations

Pregnancy: Baricitinib should be used during pregnancy only if the potential benefit justifies the potential risk for the mother and the fetus.

Renal Impairment: There are limited data for baricitinib in patients with severe renal impairment. Baricitinib is not recommended for patients who are on dialysis, have end-stage renal disease, or have acute kidney injury.

Hepatic Impairment: Baricitinib has not been studied in patients with severe hepatic impairment. Baricitinib should only be used in patients with severe hepatic impairment if the potential benefit outweighs the potential risk.

BC HCP EUA ISI 19NOV2020

Click [here](#) for resources related to Lilly's COVID-19 efforts and [here](#) for baricitinib's EUA.

Indication and Usage for OLUMIANT (baricitinib) tablets (in the United States) for RA patients

OLUMIANT® (baricitinib) 2-mg is indicated for the treatment of adult patients with moderately to severely active rheumatoid arthritis who have had an inadequate response to one or more tumor necrosis factor (TNF) antagonist therapies. Limitation of Use: Use of OLUMIANT in combination with other JAK inhibitors, biologic disease-modifying antirheumatic drugs (DMARDs), or with potent immunosuppressants such as azathioprine and cyclosporine is not recommended.

IMPORTANT SAFETY INFORMATION FOR OLUMIANT (baricitinib) TABLETS

WARNING: SERIOUS INFECTIONS, MALIGNANCY, AND THROMBOSIS

SERIOUS INFECTIONS: Patients treated with Olumiant are at risk for developing serious infections that may lead to hospitalization or death. Most patients who developed these infections were taking concomitant immunosuppressants such as methotrexate or corticosteroids. If a serious infection develops, interrupt Olumiant until the infection is controlled. Reported infections include:

- ○ Active tuberculosis (TB), which may present with pulmonary or extrapulmonary disease. Test patients for latent TB before initiating Olumiant and during therapy. If positive, start treatment for latent infection prior to Olumiant use.
- Invasive fungal infections, including candidiasis and pneumocystosis. Patients with invasive fungal infections may present with disseminated, rather than localized, disease.
- Bacterial, viral, and other infections due to opportunistic pathogens.

Carefully consider the risks and benefits of Olumiant prior to initiating therapy in patients with chronic or recurrent infection.

Closely monitor patients for the development of signs and symptoms of infection during and after treatment with Olumiant including the possible development of TB in patients who tested negative for latent TB infection prior to initiating therapy.

MALIGNANCIES: Lymphoma and other malignancies have been observed in patients treated with Olumiant.

THROMBOSIS: Thrombosis, including deep venous thrombosis (DVT) and pulmonary embolism (PE), has been observed at an increased incidence in patients treated with Olumiant compared to placebo. In addition, there were cases of arterial thrombosis. Many of these adverse events were serious and some resulted in death. Patients with symptoms of thrombosis should be promptly evaluated.

WARNINGS AND PRECAUTIONS

SERIOUS INFECTIONS: The most common serious infections reported with Olumiant included pneumonia, herpes zoster and urinary tract infection. Among opportunistic infections, tuberculosis, multidermatomal herpes zoster, esophageal candidiasis, pneumocystosis, acute histoplasmosis, cryptococcosis, cytomegalovirus and BK virus were reported with Olumiant. Some patients have presented with disseminated rather than local disease and were often taking concomitant immunosuppressants such as methotrexate or corticosteroids. Avoid Olumiant in patients with an active, serious infection, including localized infections. Consider the risks and benefits of treatment prior to initiating Olumiant in patients:

- ○ with chronic or recurrent infection
- who have been exposed to TB
- with a history of a serious or an opportunistic infection
- who have resided or traveled in areas of endemic tuberculosis or endemic mycoses; or
- with underlying conditions that may predispose them to infection.

Closely monitor patients for infections during and after Olumiant treatment. Interrupt Olumiant if a patient develops a serious infection, an opportunistic infection, or sepsis. Do not resume Olumiant until the infection is controlled.

Tuberculosis – Before initiating Olumiant evaluate and test patients for latent or active infection and treat patients with latent TB with standard antimycobacterial therapy. Olumiant should not be given to patients with active TB. Consider anti-TB therapy prior to initiating Olumiant in patients with a history of latent or active TB in whom an adequate course of treatment cannot be confirmed, and for patients with a negative test for latent TB but who have risk factors for TB infection. Monitor patients for TB during Olumiant treatment.

Viral Reactivation – Viral reactivation, including cases of herpes virus reactivation (e.g., herpes zoster), were reported in clinical studies with Olumiant. If a patient develops herpes zoster, interrupt Olumiant treatment until the episode resolves.

The impact of Olumiant on chronic viral hepatitis reactivation is unknown. Screen for viral hepatitis in accordance with clinical guidelines before initiating Olumiant.

MALIGNANCY AND LYMPHOPROLIFERATIVE DISORDERS: Malignancies were observed in Olumiant clinical studies. Consider the risks and benefits of Olumiant prior to initiating therapy in patients with a known malignancy other than a successfully treated non-melanoma skin cancer (NMSC) or when considering continuing Olumiant in patients who develop a malignancy. NMSCs were reported in patients treated with Olumiant. Periodic skin examination is recommended for patients who are at increased risk for skin cancer.

THROMBOSIS: Thrombosis, including DVT and PE, has been observed at an increased incidence in Olumiant-treated patients compared to placebo. In addition, arterial thrombosis events in the extremities have been reported in clinical studies with Olumiant. Many of these adverse events were serious and some resulted in death. There was no clear relationship between platelet count elevations and thrombotic events. Use Olumiant with caution in patients who may be at increased risk of thrombosis. If clinical features of DVT/PE or arterial thrombosis occur, evaluate patients promptly and treat appropriately.

GASTROINTESTINAL PERFORATIONS: Gastrointestinal perforations have been reported in Olumiant clinical studies, although the role of JAK inhibition in these events is not known. Use Olumiant with caution in patients who may be at increased risk for gastrointestinal perforation (e.g., patients with a history of diverticulitis). Promptly evaluate patients who present with new onset abdominal symptoms for early identification of gastrointestinal perforation.

LABORATORY ABNORMALITIES:

Neutropenia – Olumiant treatment was associated with an increased incidence of neutropenia (absolute neutrophil count [ANC] <1000 cells/mm³) compared to placebo. Avoid initiation or interrupt Olumiant treatment in patients with an ANC <1000 cells/mm³. Evaluate at baseline and thereafter according to routine patient management.

Lymphopenia – Absolute lymphocyte count (ALC) <500 cells/mm³ were reported in Olumiant clinical trials. Lymphocyte counts less than the lower limit of normal were associated with infection in patients treated with Olumiant, but not placebo. Avoid initiation or interrupt Olumiant treatment in

patients with an ALC <500 cells/mm³. Evaluate at baseline and thereafter according to routine patient management.

Anemia – Decreases in hemoglobin levels to <8 g/dL were reported in Olumiant clinical trials. Avoid initiation or interrupt Olumiant treatment in patients with hemoglobin <8 g/dL. Evaluate at baseline and thereafter according to routine patient management.

Liver Enzyme Elevations – Olumiant treatment was associated with increased incidence of liver enzyme elevation compared to placebo. Increases of ALT ≥5x upper limit of normal (ULN) and increases of AST ≥10x ULN were observed in patients in Olumiant clinical trials.

Evaluate at baseline and thereafter according to routine patient management. Promptly investigate the cause of liver enzyme elevation to identify potential cases of drug-induced liver injury. If increases in ALT or AST are observed and drug-induced liver injury is suspected, interrupt Olumiant until this diagnosis is excluded.

Lipid Elevations – Treatment with Olumiant was associated with increases in lipid parameters, including total cholesterol, low-density lipoprotein cholesterol and high-density lipoprotein cholesterol. Assess lipid parameters approximately 12 weeks following Olumiant initiation. Manage patients according to clinical guidelines for the management of hyperlipidemia.

VACCINATIONS: Avoid use of live vaccines with Olumiant. Update immunizations in agreement with current immunization guidelines prior to initiating Olumiant therapy.

HYPERSENSITIVITY: Reactions such as angioedema, urticaria, and rash that may reflect drug sensitivity have been observed in patients receiving Olumiant, including serious reactions. If a serious hypersensitivity reaction occurs, promptly discontinue Olumiant while evaluating the potential causes of the reaction.

ADVERSE REACTIONS

Most common adverse reactions include: upper respiratory tract infections (16.3%, 11.7%), nausea (2.7%, 1.6%), herpes simplex (0.8%, 0.7%) and herpes zoster (1.0%, 0.4%) for Olumiant 2 mg and placebo, respectively.

USE IN SPECIFIC POPULATIONS

PREGNANCY AND LACTATION: No information is available to support the use of Olumiant in pregnancy or lactation. Advise women not to breastfeed during treatment with Olumiant.

HEPATIC AND RENAL IMPAIRMENT: Olumiant is not recommended in patients with severe hepatic impairment or in patients with severe renal impairment.

Please click to access full [Prescribing Information](#), including [Boxed Warning about Serious infections, Malignancies, and Thrombosis](#), and [Medication Guide](#).

BA HCP ISI 09JUL2020

About OLUMIANT® (baricitinib)

OLUMIANT, a once-daily, oral JAK inhibitor was discovered by Incyte and licensed to Lilly. It is approved in the U.S. and more than 75 countries as a treatment for adults with moderate to severe rheumatoid arthritis and was recently approved in the European Union and Japan for the treatment of adult patients with moderate to severe atopic dermatitis who are candidates for systemic therapy. Olumiant was recently approved in Japan for the treatment of pneumonia associated with COVID-19 in hospitalized adult patients. The U.S. FDA-approved labeling for Olumiant includes a Boxed Warning for Serious Infections, Malignancy, and Thrombosis. See the full Prescribing Information [here](#). Baricitinib is also being investigated in alopecia areata (AA), juvenile idiopathic arthritis (JIA) and systemic lupus erythematosus (SLE).

In December 2009, Lilly and Incyte announced an exclusive worldwide license and collaboration agreement for the development and commercialization of baricitinib and certain follow-on compounds for patients with inflammatory and autoimmune diseases.

About Lilly's COVID-19 Efforts

Lilly is bringing the full force of its scientific and medical expertise to attack the coronavirus pandemic around the world. Existing Lilly medicines are being studied to understand their potential in treating complications of COVID-19, and the company is collaborating with partner companies to discover and develop novel antibody treatments for COVID-19. Click [here](#) for resources related to Lilly's COVID-19 efforts.

About Eli Lilly and Company

Lilly is a global health care leader that unites caring with discovery to create medicines that make life better for people around the world. We were founded more than a century ago by a man committed to creating high-quality medicines that meet real needs, and today we remain true to that mission in all our work. Across the globe, Lilly employees work to discover and bring life-changing medicines to those who need them, improve the understanding and management of disease, and give back to communities through philanthropy and volunteerism. To learn more about Lilly, please visit us at www.lilly.com and www.lilly.com/news. P-LLY

About Direct Relief

Direct Relief is a humanitarian aid organization, active in all 50 states and more than 80 countries, with a mission to improve the health and lives of people affected by poverty or emergencies – without regard to politics, religion, or ability to pay.

Lilly Cautionary Statement Regarding Forward-Looking Statements

This press release contains forward-looking statements (as that term is defined in the Private Securities Litigation Reform Act of 1995) about OLUMIANT (baricitinib) as a potential treatment for patients with COVID-19 and reflects Lilly's current beliefs and expectations. However, as with any pharmaceutical product, there are substantial risks and uncertainties in the process of research, development and commercialization. Among other things, there can be no guarantee that planned or ongoing studies will be completed as planned, that future study results will be consistent with the results to date, that OLUMIANT will receive additional regulatory approvals or authorizations or be commercially successful, that OLUMIANT will be safe and effective as a treatment or successful preventative therapy for COVID-19 or that Lilly's licensing arrangements will provide an adequate supply of OLUMIANT. For further discussion of these and other risks and uncertainties, see Lilly's most recent Form 10-K and Form 10-Q filings with the United States Securities and Exchange Commission. Except as required by law, Lilly undertakes no duty to update forward-looking statements to reflect events after the date of this release.

#

Refer to: Kristen Basu; basu_kristen_porter@lilly.com; 317-447-2199 (Media)

Kevin Hern; hern_kevin_r@lilly.com; 317-277-1838 (Investors)

Lilly

View original content to download multimedia:<http://www.prnewswire.com/news-releases/lilly-accelerating-baricitinibs-availability-in-india-following-receipt-of-permission-for-restricted-emergency-use-as-a-covid-19-therapy-via-donations-and-licensing-agreements-301282984.html>

SOURCE Eli Lilly and Company