



Lilly to supply 614,000 additional doses of bamlanivimab and etesevimab to the U.S. Government for the treatment or post-exposure prevention of COVID-19

November 2, 2021

INDIANAPOLIS, Nov. 2, 2021 /PRNewswire/ -- Eli Lilly and Company (NYSE: LLY) today announced an additional purchase by the U.S. government for bamlanivimab with etesevimab for administration together. This neutralizing antibody therapy is authorized for emergency use for the treatment of mild to moderate COVID-19 or for post-exposure prophylaxis of COVID-19 in certain individuals. As part of the agreement, Lilly will supply 614,000 doses of bamlanivimab with etesevimab no later than January 31, 2022 for a total of \$1.29 billion. A minimum of 400,000 doses will be supplied no later than December 31, 2021.

The estimated impact of the new purchase to the 2021 guidance provided in Lilly's earnings release dated October 26, 2021 is \$840 million of additional revenue and approximately 25 cents of additional earnings per share. The expected impact in 2022 will be reflected in the company's financial guidance to be provided in December 2021.

For more information about the use of bamlanivimab with etesevimab to treat COVID-19, click [here](#) or contact Lilly's 24-hour support line at 1-855-LillyC19 (1-855-545-5921). For media resources, including product images and fact sheets, please click [here](#).

This purchase has been supported in whole or in part with federal funds from the Department of Health and Human Services; Office of the Assistant Secretary for Preparedness and Response; Biomedical Advanced Research and Development Authority (BARDA), under Contract No. W911QY21D0012.

Important Information about bamlanivimab and etesevimab together

Bamlanivimab and etesevimab together have not been approved by the FDA for any use. It is not known if bamlanivimab and etesevimab together are safe and effective for the treatment or post-exposure prophylaxis of COVID-19.

Bamlanivimab and etesevimab together are authorized under Emergency Use Authorization only for the duration of the declaration that circumstances exist justifying the authorization of the emergency use under Section 564(b)(1) of the Act, 21 U.S.C § 360bbb-3(b)(1), unless the authorization is terminated or revoked sooner.

Healthcare providers should review the Fact Sheet for information on the authorized use of bamlanivimab and etesevimab together and mandatory requirements of the EUA. Please see the [FDA Letter of Authorization](#), [Fact Sheet for Healthcare Providers](#), and Fact Sheet for Patients, Parents and Caregivers ([English](#)) ([Spanish](#)) for bamlanivimab and etesevimab together.

Authorized Use and Important Safety Information

TREATMENT

Bamlanivimab and etesevimab together are authorized for the treatment of mild to moderate coronavirus disease 2019 (COVID-19) in adults and pediatric patients (12 years of age and older weighing at least 40 kg) with positive results of direct SARS-CoV-2 viral testing, and who are at high risk for progression to severe COVID-19, including hospitalization or death.

Limitations of Authorized Use: Treatment

Combined Frequency of Variants Resistant to Bamlanivimab and Etesevimab

- Bamlanivimab and etesevimab are not authorized for use in states, territories, and U.S. jurisdictions in which the combined frequency of variants resistant to bamlanivimab and etesevimab exceeds 5%.ⁱ
 - A list of states, territories, and U.S. jurisdictions in which bamlanivimab and etesevimab are and are not currently authorized is available on the following [FDA website](#).

Use in Patients Who Are Hospitalized or Who Require Oxygen Due to COVID-19

Bamlanivimab and etesevimab together are not authorized for use in patients:

- who are hospitalized due to COVID-19, OR
- who require oxygen therapy due to COVID-19, OR
- who require an increase in baseline oxygen flow rate due to COVID-19 in those on chronic oxygen therapy due to underlying non-COVID-19 related comorbidity.

Treatment with bamlanivimab and etesevimab has not been studied in patients hospitalized due to COVID-19. Monoclonal antibodies, such as bamlanivimab and etesevimab, may be associated with worse clinical outcomes when administered to hospitalized patients with COVID-19 requiring high-flow oxygen or mechanical ventilation.

POST-EXPOSURE PROPHYLAXIS

Bamlanivimab and etesevimab administered together are authorized in adults and pediatric individuals (12 years of age or older weighing at least 40 kg) for post-exposure prophylaxis of COVID-19 in individuals who are at high risk for progression to severe COVID-19, including hospitalization or death, and are:

- not fully vaccinatedⁱⁱ or who are not expected to mount an adequate immune response to complete SARS-CoV-2 vaccination (for example, individuals with immunocompromising conditions including those taking immunosuppressive medicationsⁱⁱⁱ) **AND**
 - have been exposed to an individual infected with SARS-CoV-2 consistent with close contact criteria per Centers for Disease Control and Prevention (CDC)^{iv} **OR**
 - who are at high risk of exposure to an individual infected with SARS-CoV-2 because of occurrence of SARS-CoV-2 infection in other individuals in the same institutional setting (for example, nursing homes, prisons).

Limitations of Authorized Use: Post-Exposure Prophylaxis

- Bamlanivimab and etesevimab are not authorized for use in states, territories, and U.S. jurisdictions in which the combined frequency of variants resistant to bamlanivimab and etesevimab exceeds 5%.ⁱ
 - A list of states, territories, and U.S. jurisdictions in which bamlanivimab and etesevimab are and are not currently authorized is available on the following [FDA website](#).
- Post-exposure prophylaxis with bamlanivimab and etesevimab is not a substitute for vaccination against COVID-19.
- Bamlanivimab and etesevimab together are not authorized for pre-exposure prophylaxis for prevention of COVID-19.

Important Safety Information

There are limited clinical data available for bamlanivimab and etesevimab. Serious and unexpected adverse events may occur that have not been previously reported with the use of bamlanivimab and etesevimab together.

Warnings

Hypersensitivity Including Anaphylaxis and Infusion-Related Reactions

Serious hypersensitivity reactions, including anaphylaxis, have been observed with administration of bamlanivimab and etesevimab. If signs and symptoms of a clinically significant hypersensitivity reaction or anaphylaxis occur, immediately discontinue administration and initiate appropriate medications and/or supportive therapy.

Infusion-related reactions, occurring during the infusion and up to 24 hours after infusion, have been observed with administration of bamlanivimab and etesevimab together. These reactions may be severe or life threatening. Signs and symptoms of infusion-related reactions may include:

- fever, difficulty breathing, reduced oxygen saturation, chills, fatigue, arrhythmia (e.g. atrial fibrillation, sinus tachycardia, bradycardia), chest pain or discomfort, weakness, altered mental status, nausea, headache, bronchospasm, hypotension, hypertension, angioedema, throat irritation, rash including urticaria, pruritus, myalgia, vasovagal reactions (e.g. presyncope, syncope), dizziness, and diaphoresis.

Consider slowing or stopping the infusion and administer appropriate medications and/or supportive care if an infusion-related reaction occurs.

Hypersensitivity reactions occurring more than 24 hours after the infusion have also been reported with the use of bamlanivimab and etesevimab under Emergency Use Authorization.

Clinical Worsening After Receiving Bamlanivimab and Etesevimab Administration

Clinical worsening of COVID-19 after administration of bamlanivimab and etesevimab together has been reported and may include signs or symptoms of fever, hypoxia or increased respiratory difficulty, arrhythmia (e.g., atrial fibrillation, sinus tachycardia, bradycardia), fatigue, and altered mental status. Some of these events required hospitalization. It is not known if these events were related to bamlanivimab and etesevimab use or were due to progression of COVID-19.

Limitations of Benefit and Potential Risk in Patients with Severe COVID-19

Treatment with bamlanivimab and etesevimab has not been studied in patients hospitalized due to COVID-19. Monoclonal antibodies, such as bamlanivimab and etesevimab, may be associated with worse clinical outcomes when administered to hospitalized patients with COVID-19 requiring high-flow oxygen or mechanical ventilation. See Limitations of Authorized Use.

Adverse Reactions

Adverse reactions observed in those who have received bamlanivimab and etesevimab are anaphylaxis (n=1, 0.07%) and infusion-related reactions (n=16, 1.1%). The most common treatment-emergent adverse events included nausea, dizziness, and pruritus. No treatment-emergent events occurred in more than 1% of participants and rates were comparable to placebo.

Use in Specific Populations

Pregnancy

There are insufficient data to evaluate a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. Bamlanivimab and etesevimab should only be used during pregnancy if the potential benefit outweighs the potential risk for the mother and the fetus.

Breastfeeding

There are no available data on the presence of bamlanivimab or etesevimab in human or animal milk, the effects on the breastfed infant, or the effects on milk production. Breastfeeding individuals with COVID-19 should follow practices according to clinical guidelines to avoid exposing the infant to COVID-19.

BM ET HCP EUA ISI 27AUG2021

About bamlanivimab and etesevimab

Bamlanivimab is a recombinant, neutralizing human IgG1 monoclonal antibody (mAb) directed against the spike protein of SARS-CoV-2. It was designed to block viral attachment and entry into human cells, thus neutralizing the virus. Bamlanivimab emerged from the collaboration

between Lilly and AbCellera to create antibody therapies for the prevention and treatment of COVID-19. Lilly scientists rapidly developed the antibody in less than three months after it was discovered by AbCellera and the scientists at the National Institute of Allergy and Infectious Diseases (NIAID) Vaccine Research Center. Bamlanivimab was identified from a blood sample taken from one of the first U.S. patients who recovered from COVID-19.

Etesevimab (LY-CoV016, also known as JS016) is a recombinant fully human monoclonal neutralizing antibody, which specifically binds to the SARS-CoV-2 surface spike protein receptor binding domain with high affinity and can block the binding of the virus to the ACE2 host cell surface receptor. Point mutations were introduced into the native human IgG1 antibody to mitigate effector function. Lilly licensed etesevimab from Junshi Biosciences after it was jointly developed by Junshi Biosciences and the Institute of Microbiology, Chinese Academy of Science (IMCAS). Junshi Biosciences leads development in Greater China, while Lilly leads development in the rest of the world.

Results from a Phase 2/3 study in people recently diagnosed with COVID-19 in the ambulatory setting (BLAZE-1, [NCT04427501](#)) were published in the *New England Journal of Medicine*. Results from a Phase 3 study of bamlanivimab in residents and staff at long-term care facilities (BLAZE-2, [NCT04497987](#)) were published in the *Journal of American Medical Association* (JAMA). A Phase 2 study assessing the efficacy and safety of bamlanivimab alone, and bamlanivimab with other neutralizing antibodies versus placebo for the treatment of symptomatic low-risk COVID-19 in the outpatient setting (BLAZE-4, [NCT04634409](#)) has completed enrollment.

About Lilly's COVID-19 Efforts

Lilly is bringing the full force of its scientific and medical expertise to attack the coronavirus pandemic around the world. Existing Lilly medicines are being studied to understand their potential in treating complications of COVID-19, and the company is collaborating with partner companies to discover and develop novel antibody therapies for COVID-19. Click [here](#) for resources related to Lilly's COVID-19 efforts.

About Eli Lilly and Company

Lilly is a global health care leader that unites caring with discovery to create medicines that make life better for people around the world. We were founded more than a century ago by a man committed to creating high-quality medicines that meet real needs, and today we remain true to that mission in all our work. Across the globe, Lilly employees work to discover and bring life-changing medicines to those who need them, improve the understanding and management of disease, and give back to communities through philanthropy and volunteerism. To learn more about Lilly, please visit us at [www.lilly.com](#) and [www.lilly.com/news](#). P-LLY

Lilly Cautionary Statement Regarding Forward-Looking Statements

This press release contains forward-looking statements (as that term is defined in the Private Securities Litigation Reform Act of 1995) about bamlanivimab (LY-CoV555) and etesevimab (LY-CoV016) together as a potential therapy for patients with COVID-19, the supply, distribution and contracts with governments relating to these therapies, and Lilly's development plans, and reflects Lilly's current beliefs and expectations. However, as with any such undertaking, there are substantial risks and uncertainties in the process of drug research, development and commercialization. Among other things, there can be no guarantee that future study results will be consistent with the results to date, that bamlanivimab and etesevimab together will prove to be a safe and effective treatment or successful preventative therapy for COVID-19, that bamlanivimab and etesevimab together will receive regulatory approvals or additional authorizations, that patients will volunteer to participate in clinical trials or achieve positive outcomes, that Lilly will obtain any additional purchase orders or supply contracts, or that Lilly can provide an adequate supply of bamlanivimab and etesevimab together in all circumstances. For a further discussion of these and other risks and uncertainties that could cause actual results to differ from Lilly's expectations, please see Lilly's most recent Forms 10-K and 10-Q filed with the U.S. Securities and Exchange Commission. Lilly undertakes no duty to update forward-looking statements.

ⁱ FDA will make this determination considering current variant frequency data (available at: <https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/variant-proportions.html>), trends in variant frequency over time, the precision of the estimates and information regarding emerging variants of concern. FDA will update the list of states, territories, and US jurisdictions in which bamlanivimab and etesevimab are and are not currently authorized as new data and information becomes available. Healthcare providers should refer to the FDA website regularly for updates. Healthcare providers should refer to the FDA website regularly for updates.

ⁱⁱ Individuals are considered to be fully vaccinated 2 weeks after their second vaccine dose in a 2-dose series (such as the Pfizer or Moderna vaccines), or 2 weeks after a single-dose vaccine (such as Johnson & Johnson's Janssen vaccine). See this website for more details: <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/fully-vaccinated.html#vaccinated>.

ⁱⁱⁱ See this website for more details: <https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/fully-vaccinated-people.html>.

^{iv} Close contact with an infected individual is defined as: being within 6 feet for a total of 15 minutes or more, providing care at home to someone who is sick, having direct physical contact with the person (hugging or kissing, for example), sharing eating or drinking utensils, or being exposed to respiratory droplets from an infected person (sneezing or coughing, for example). See this website for additional details: <https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/quarantine.html>.

Refer to: Allison Howell; howell_allison@lilly.com; 317-655-1106 (Media)
Kevin Hern; hern_kevin_r@lilly.com; 317-277-1838 (Investors)

Lilly

View original content to download multimedia: <https://www.prnewswire.com/news-releases/lilly-to-supply-614-000-additional-doses-of-bamlanivimab-and-etesevimab-to-the-us-government-for-the-treatment-or-post-exposure-prevention-of-covid-19--301413637.html>

SOURCE Eli Lilly and Company