Lilly Announces Details of Presentations at 2023 American Association for Cancer Research (AACR) Annual Meeting

March 14, 2023

INDIANAPOLIS, March 14, 2023 /PRNewswire/ -- Eli Lilly and Company (NYSE: LLY) today announced that data from its oncology portfolio will be presented at the American Association for Cancer Research (AACR) Annual Meeting in Orlando, April 14 – 19, 2023. The presentations include the first clinical data from the Phase 1 studies of LY3537982 (KRAS G12C inhibitor) and LY3410738 (IDH inhibitor) and the Phase 2 study of Verzenio® (abemaciclib; CDK4/6 inhibitor) in metastatic treatment refractory castration-resistant prostate cancer. In addition, preclinical characterization data on Jaypirca's™ (pirtobrutinib; BTK inhibitor) binding mechanism will also be presented.

A list of the presentations, along with their viewing details, are shared below.

Medicine	***Abstract Title***	***Presentation Details***
LY3537982	A first-in-human phase 1 study of LY3537982, a highly selective and potent KRAS G12C inhibitor in patients with KRAS G12C-mutant advanced solid tumors	**Abstract Number:** CT028 **Session Title**: Targeting the KRAS Pathway in the Clinic **Session Type**: Oral **Session Date / Time:** April 17 at 2:30 PM – 4:30 PM ET
LY3410738	A first-in-human phase 1 study of LY3410738, a covalent inhibitor of mutant IDH, in advanced myeloid malignancies	**Abstract Number:** CT026 **Session Title:** Novel Clinical Trials for Hematological Malignancies **Session Type:** Oral **Session Date / Time**: April 16 at 3:00 PM – 5:00 PM ET
LY3410738	A first-in-human phase 1 study of LY3410738, a covalent inhibitor of mutant IDH, in advanced IDH-mutant cholangiocarcinoma and other solid tumors	**Abstract Number**: CT098 **Session Title**: First-in-Human Phase I Clinical Trials 1 **Session Type**: Poster **Session Date / Time:** April 17 at 1:30 PM – 5:00 PM ET **Location:** Poster Section 45
Verzenio® (abemaciclib)	CYCLONE 1: A Phase 2 Study of Abemaciclib in Metastatic Castration-Resistant Prostate Cancer Patients Previously Treated with a Novel Hormonal Agent and Taxane-based Chemotherapy	**Abstract Number:** CT159 **Session Title:** Phase II Clinical Trials 1 **Session Type**: Poster **Session Date / Time:** April 17 at 1:30 PM – 5:00 PM ET **Location**: Poster Section 47
Jaypirca™(pirtobrutinib)	Unique pharmacodynamic properties conferred by differential binding to BTK, pirtobrutinib vs covalent inhibitors	**Abstract Number:** 2780 **Session Title**: Pharmacokinetics, Pharmacodynamics, and Molecular Pharmacology **Session Type**: Poster **Session Date / Time**: April 17 at 1:30 PM – 5:00 PM ET **Location:** Poster Section 18

INDICATIONS FOR VERZENIO®

VERZENIO® is a kinase inhibitor indicated:

- in combination with endocrine therapy (tamoxifen or an aromatase inhibitor) for the adjuvant treatment of adult patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, node-positive, early breast cancer at high risk of recurrence.
- in combination with an aromatase inhibitor as initial endocrine-based therapy for the treatment of adult patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer.
- in combination with fulvestrant for the treatment of adult patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer with disease progression following endocrine therapy.
- as monotherapy for the treatment of adult patients with HR-positive, HER2-negative advanced or metastatic breast cancer

with disease progression following endocrine therapy and prior chemotherapy in the metastatic setting.

IMPORTANT SAFETY INFORMATION FOR VERZENIO (abemaciclib)
Severe **diarrhea** associated with dehydration and infection occurred in patients treated with Verzenio. Across four clinical trials in 3691 patients, diarrhea occurred in 81 to 90% of patients who received Verzenio. Grade 3 diarrhea occurred in 8 to 20% of patients receiving Verzenio. Most patients experienced diarrhea during the first month of Verzenio treatment. The median time to onset of the first diarrhea event ranged from 6 to 8 days; and the median duration of Grade 2 and Grade 3 diarrhea ranged from 6 to 11 days and 5 to 8 days, respectively. Across trials, 19 to 26% of patients with diarrhea required a Verzenio dose interruption and 13 to 23% required a dose reduction.

Instruct patients to start antidiarrheal therapy, such as loperamide, at the first sign of loose stools, increase oral fluids, and notify their healthcare provider for further instructions and appropriate follow-up. For Grade 3 or 4 diarrhea, or diarrhea that requires hospitalization, discontinue Verzenio until toxicity resolves to ≤Grade 1, and then resume Verzenio at the next lower dose.

Neutropenia, including febrile neutropenia and fatal neutropenic sepsis, occurred in patients treated with Verzenio. Across four clinical trials in 3691 patients, neutropenia occurred in 37 to 46% of patients receiving Verzenio. A Grade ≥3 decrease in neutrophil count (based on laboratory findings) occurred in 19 to 32% of patients receiving Verzenio. Across trials, the median time to first episode of Grade ≥3 neutropenia ranged from 29 to 33 days, and the median duration of Grade ≥3 neutropenia ranged from 11 to 16 days. Febrile neutropenia has been reported in <1% of patients exposed to Verzenio across trials. Two deaths due to neutropenic sepsis were observed in MONARCH 2. Inform patients to promptly report any episodes of fever to their healthcare provider.

Monitor complete blood counts prior to the start of Verzenio therapy, every 2 weeks for the first 2 months, monthly for the next 2 months, and as clinically indicated. Dose interruption, dose reduction, or delay in starting treatment cycles is recommended for patients who develop Grade 3 or 4 neutropenia.

Severe, life-threatening, or fatal **interstitial lung disease (ILD) or pneumonitis** can occur in patients treated with Verzenio and other CDK4/6 inhibitors. In Verzenio-treated patients in EBC (monarchE), 3% of patients experienced ILD or pneumonitis of any grade: 0.4% were Grade 3 or 4 and there was one fatality (0.1%). In Verzenio-treated patients in MBC (MONARCH 1, MONARCH 2, MONARCH 3), 3.3% of Verzenio-treated patients had ILD or pneumonitis of any grade: 0.6% had Grade 3 or 4, and 0.4% had fatal outcomes. Additional cases of ILD or pneumonitis have been observed in the postmarketing setting, with fatalities reported.

Monitor patients for pulmonary symptoms indicative of ILD or pneumonitis. Symptoms may include hypoxia, cough, dyspnea, or interstitial infiltrates on radiologic exams. Infectious, neoplastic, and other causes for such symptoms should be excluded by means of appropriate investigations. Dose interruption or dose reduction is recommended in patients who develop persistent or recurrent Grade 2 ILD or pneumonitis. Permanently discontinue Verzenio in all patients with Grade 3 or 4 ILD or pneumonitis.

Grade ≥3 increases in alanine aminotransferase (ALT) (2 to 6%) and **aspartate aminotransferase (AST)** (2 to 3%) were reported in patients receiving Verzenio. Across three clinical trials in 3559 patients (monarchE, MONARCH 2, MONARCH 3), the median time to onset of Grade ≥3 ALT increases ranged from 57 to 87 days and the median time to resolution to Grade <3 was 13 to 14 days. The median time to onset of Grade ≥3 AST increases ranged from 71 to 185 days and the median time to resolution to Grade <3 ranged from 11 to 15 days.

Monitor liver function tests (LFTs) prior to the start of Verzenio therapy, every 2 weeks for the first 2 months, monthly for the next 2 months, and as clinically indicated. Dose interruption, dose reduction, dose discontinuation, or delay in starting treatment cycles is recommended for patients who develop persistent or recurrent Grade 2, or any Grade 3 or 4 hepatic transaminase elevation.

Venous thromboembolic events (VTE) were reported in 2 to 5% of patients across three clinical trials in 3559 patients treated with Verzenio (monarchE, MONARCH 2, MONARCH 3). VTE included deep vein thrombosis, pulmonary embolism, pelvic venous thrombosis, cerebral venous sinus thrombosis, subclavian and axillary vein thrombosis, and inferior vena cava thrombosis. In clinical trials, deaths due to VTE have been reported in patients treated with Verzenio.

Verzenio has not been studied in patients with early breast cancer who had a history of VTE. Monitor patients for signs and symptoms of venous thrombosis and pulmonary embolism and treat as medically appropriate. Dose interruption is recommended for EBC patients with any grade VTE and for MBC patients with a Grade 3 or 4 VTE.

Verzenio can cause **fetal harm** when administered to a pregnant woman, based on findings from animal studies and the mechanism of action. In animal reproduction studies, administration of abemaciclib to pregnant rats during the period of organogenesis caused teratogenicity and decreased fetal weight at maternal exposures that were similar to the human clinical exposure based on area under the curve (AUC) at the maximum recommended human dose. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with Verzenio and for 3 weeks after the last dose. Based on findings in animals, Verzenio may impair fertility in males of reproductive potential. There are no data on the presence of Verzenio in human milk or its effects on the breastfed child or on milk production. Advise lactating women not to breastfeed during Verzenio treatment and for at least 3 weeks after the last dose because of the potential for serious adverse reactions in breastfed infants.

The **most common adverse reactions (all grades, ≥10%)** observed in **monarchE for Verzenio plus tamoxifen or an aromatase inhibitor vs tamoxifen or an aromatase inhibitor, with a difference between arms of ≥2%,** were diarrhea (84% vs 9%), infections (51% vs 39%), neutropenia (46% vs 6%), fatigue (41% vs 18%), leukopenia (38% vs 7%), nausea (30% vs 9%), anemia (24% vs 4%), headache (20% vs 15%), vomiting (18% vs 4.6%), stomatitis (14% vs 5%), lymphopenia (14% vs 3%), thrombocytopenia (13% vs 2%), decreased appetite (12% vs 2.4%), ALT increased (12% vs 6%), AST increased (12% vs 5%), dizziness (11% vs 7%), rash (11% vs 4.5%), and alopecia (11% vs 2.7 %).

The **most frequently reported ≥5% Grade 3 or 4 adverse reaction** that occurred in the Verzenio arm vs the tamoxifen or an aromatase inhibitor arm of monarchE were neutropenia (19.6% vs 1%), leukopenia (11% vs <1%), diarrhea (8% vs 0.2%), and lymphopenia (5% vs <1%).

Lab abnormalities (all grades; Grade 3 or 4) for **monarchE in ≥10% for Verzenio plus tamoxifen or an aromatase inhibitor with a difference between arms of ≥2%** were increased serum creatinine (99% vs 91%; .5% vs <.1%), decreased white blood cells (89% vs 28%; 19.1% vs 1.1%), decreased neutrophil count (84% vs 23%; 18.7% vs 1.9%), anemia (68% vs 17%; 1% vs .1%), decreased lymphocyte count (59% vs 24%; 13.2 % vs 2.5%), decreased platelet count (37% vs 10%; .9% vs .2%), increased ALT (37% vs 24%; 2.6% vs 1.2%), increased AST (31% vs 18%; 1.6% vs .9%),

and hypokalemia (11% vs 3.8%; 1.3% vs 0.2%).

The **most common adverse reactions (all grades, ≥10%)** observed in **MONARCH 3 for Verzenio plus anastrozole or letrozole vs anastrozole or letrozole, with a difference between arms of ≥2%,** were diarrhea (81% vs 30%), fatigue (40% vs 32%), neutropenia (41% vs 2%), infections (39% vs 29%), nausea (39% vs 20%), abdominal pain (29% vs 12%), vomiting (28% vs 12%), anemia (28% vs 5%), alopecia (27% vs 11%), decreased appetite (24% vs 9%), leukopenia (21% vs 2%), creatinine increased (19% vs 4%), constipation (16% vs 12%), ALT increased (16% vs 7%), AST increased (15% vs 7%), rash (14% vs 5%), pruritus (13% vs 9%), cough (13% vs 9%), dyspnea (12% vs 6%), dizziness (11% vs 9%), weight decreased (10% vs 3.1%), influenza-like illness (10% vs 8%), and thrombocytopenia (10% vs 2%).

The **most frequently reported ≥5% Grade 3 or 4 adverse reactions** that occurred in the Verzenio arm vs the placebo arm of **MONARCH 3** were neutropenia (22% vs 1%), diarrhea (9% vs 1.2%), leukopenia (7% vs <1%)), increased ALT (6% vs 2%), and anemia (6% vs 1%).

Lab abnormalities (all grades; Grade 3 or 4) for **MONARCH 3 in ≥10% for Verzenio plus anastrozole or letrozole with a difference between arms of ≥2%** were increased serum creatinine (98% vs 84%; 2.2% vs 0%), decreased white blood cells (82% vs 27%; 13% vs 0.6%), anemia (82% vs 28%; 1.6% vs 0%), decreased neutrophil count (80% vs 21%; 21.9% vs 2.6%), decreased lymphocyte count (53% vs 26%; 7.6% vs 1.9%), decreased platelet count (36% vs 12%; 1.9% vs 0.6%), increased ALT (48% vs 25%; 6.6% vs 1.9%), and increased AST (37% vs 23%; 3.8% vs 0.6%).

The **most common adverse reactions (all grades, ≥10%)** observed in **MONARCH 2 for Verzenio plus fulvestrant vs fulvestrant, with a difference between arms of ≥2%,** were diarrhea (86% vs 25%), neutropenia (46% vs 4%), fatigue (46% vs 32%), nausea (45% vs 23%), infections (43% vs 25%), abdominal pain (35% vs 16%), anemia (29% vs 4%), leukopenia (28% vs 2%), decreased appetite (27% vs 12%), vomiting (26% vs 10%), headache (20% vs 15%), dysgeusia (18% vs 2.7%), thrombocytopenia (16% vs 3%), alopecia (16% vs 1.8%), stomatitis (15% vs 10%), ALT increased (13% vs 5%), pruritus (13% vs 6%), cough (13% vs 11%), dizziness (12% vs 6%), AST increased (12% vs 7%), peripheral edema (12% vs 7%), creatinine increased (12% vs <1%), rash (11% vs 4.5%), pyrexia (11% vs 6%), and weight decreased (10% vs 2.2%).

The **most frequently reported ≥5% Grade 3 or 4 adverse reactions** that occurred in the Verzenio arm vs the placebo arm of **MONARCH 2** were neutropenia (25% vs 1%), diarrhea (13% vs 0.4%), leukopenia (9% vs 0%), anemia (7% vs 1%), and infections (5.7% vs 3.5%).

Lab abnormalities (all grades; Grade 3 or 4) for **MONARCH 2 in ≥10% for Verzenio plus fulvestrant with a difference between arms of ≥2%** were increased serum creatinine (98% vs 74%; 1.2% vs 0%), decreased white blood cells (90% vs 33%; 23.7% vs .9%), decreased neutrophil count (87% vs 30%; 32.5% vs 4.2%), anemia (84% vs 34%; 2.6% vs .5%), decreased lymphocyte count (63% vs 32%; 12.2% vs 1.8%), decreased platelet count (53% vs 15%; 2.1% vs 0%), increased ALT (41% vs 32%; 4.6% vs 1.4%), and increased AST (37% vs 25%; 3.9% vs 4.2%).

The **most common adverse reactions (all grades, ≥10%)** observed in **MONARCH 1** with Verzenio were diarrhea (90%), fatigue (65%), nausea (64%), decreased appetite (45%), abdominal pain (39%), neutropenia (37%), vomiting (35%), infections (31%), anemia (25%), thrombocytopenia (20%), headache (20%), cough (19%), constipation (17%), leukopenia (17%), arthralgia (15%), dry mouth (14%), weight decreased (14%), stomatitis (14%), creatinine increased (13%), alopecia (12%), dysgeusia (12%), pyrexia (11%), dizziness (11%), and dehydration (10%).

The **most frequently reported ≥5% Grade 3 or 4 adverse reactions** from **MONARCH 1** with Verzenio were diarrhea (20%), neutropenia (24%), fatigue (13%), and leukopenia (5%).

Lab abnormalities (all grades; Grade 3 or 4) for MONARCH 1 with Verzenio were increased serum creatinine (99%; .8%), decreased white blood cells (91%; 28%), decreased neutrophil count (88%; 26.6%), anemia (69%; 0%), decreased lymphocyte count (42%; 13.8%), decreased platelet count (41%; 2.3%), increased ALT (31%; 3.1%), and increased AST (30%; 3.8%).

Strong and moderate CYP3A inhibitors increased the exposure of abemaciclib plus its active metabolites to a clinically meaningful extent and may lead to increased toxicity. Avoid concomitant use of ketoconazole. Ketoconazole is predicted to increase the AUC of abemaciclib by up to 16-fold. In patients with recommended starting doses of 200 mg twice daily or 150 mg twice daily, reduce the Verzenio dose to 100 mg twice daily with concomitant use of strong CYP3A inhibitors other than ketoconazole. In patients who have had a dose reduction to 100 mg twice daily due to adverse reactions, further reduce the Verzenio dose to 50 mg twice daily with concomitant use of strong CYP3A inhibitors. If a patient taking Verzenio discontinues a strong CYP3A inhibitor, increase the Verzenio dose (after 3 to 5 half-lives of the inhibitor) to the dose that was used before starting the inhibitor. With concomitant use of moderate CYP3A inhibitors, monitor for adverse reactions and consider reducing the Verzenio dose in 50 mg decrements. Patients should avoid grapefruit products.

Avoid concomitant use of strong or moderate CYP3A inducers and consider alternative agents. Coadministration of strong or moderate CYP3A inducers decreased the plasma concentrations of abemaciclib plus its active metabolites and may lead to reduced activity.

With severe hepatic impairment (Child-Pugh C), reduce the Verzenio dosing frequency to once daily. The pharmacokinetics of Verzenio in patients with **severe renal impairment** (CLcr <30 mL/min), end stage renal disease, or in patients on dialysis is unknown. No dosage adjustments are necessary in patients with mild or moderate hepatic (Child-Pugh A or B) and/or renal impairment (CLcr ≥30-89 mL/min).

AL HCP ISI 12OCT2021

Please see full Prescribing Information and Patient Information for Verzenio.

INDICATION FOR JAYPIRCA:

Jaypirca is a kinase inhibitor indicated for the treatment of adult patients with relapsed or refractory (R/R) mantle cell lymphoma (MCL) after at least two lines of systemic therapy, including a BTK inhibitor.

This indication is approved under accelerated approval based on response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

IMPORTANT SAFETY INFORMATION FOR JAYPIRCA ™(pirtobrutinib)
Infections: Fatal and serious infections (including bacterial, viral, or fungal) and opportunistic infections have occurred in patients treated with Jaypirca. In the clinical trial, Grade ≥3 infections occurred in 17% of 583 patients with hematologic malignancies, most commonly pneumonia (9%); fatal infections occurred in 4.1% of patients. Sepsis (4.5%) and febrile neutropenia (2.9%) occurred. Opportunistic infections after Jaypirca treatment

included, but are not limited to, *Pneumocystis jirovecii* pneumonia and fungal infection. Consider prophylaxis, including vaccinations and antimicrobial prophylaxis, in patients at increased risk for infection, including opportunistic infections. Monitor patients for signs and symptoms, evaluate promptly, and treat appropriately. Based on severity, reduce dose, temporarily withhold, or permanently discontinue Jaypirca.

Hemorrhage: Fatal and serious hemorrhage has occurred with Jaypirca. Major hemorrhage (Grade ≥3 bleeding or any central nervous system bleeding) occurred in 2.4% of 583 patients with hematologic malignancies treated with Jaypirca, including gastrointestinal hemorrhage; fatal hemorrhage occurred in 0.2% of patients. Bleeding of any grade, excluding bruising and petechiae, occurred in 14% of patients. Major hemorrhage occurred in patients taking Jaypirca with (0.7%) and without (1.7%) antithrombotic agents. Consider risks/benefits of co-administering antithrombotic agents with Jaypirca. Monitor patients for signs of bleeding. Based on severity, reduce dose, temporarily withhold, or permanently discontinue Jaypirca. Consider benefit/risk of withholding Jaypirca 3-7 days pre- and post-surgery depending on type of surgery and bleeding risk.

Cytopenias: Grade 3 or 4 cytopenias, including neutropenia (24%), anemia (11%), and thrombocytopenia (11%), have developed in patients with hematologic malignancies treated with Jaypirca. In a clinical trial, Grade 4 neutropenia (13%) and Grade 4 thrombocytopenia (5%) developed. Monitor complete blood counts regularly during treatment. Based on severity, reduce dose, temporarily withhold, or permanently discontinue Jaypirca.

Atrial Fibrillation and Atrial Flutter: Atrial fibrillation or flutter were reported in 2.7% of patients, with Grade 3 or 4 atrial fibrillation or flutter reported in 1% of 583 patients with hematologic malignancies treated with Jaypirca. Patients with cardiac risk factors such as hypertension or previous arrhythmias may be at increased risk. Monitor for signs and symptoms of arrhythmias (e.g., palpitations, dizziness, syncope, dyspnea) and manage appropriately. Based on severity, reduce dose, temporarily withhold, or permanently discontinue Jaypirca.

Second Primary Malignancies: Second primary malignancies, including non-skin carcinomas, developed in 6% of 583 patients with hematologic malignancies treated with Jaypirca monotherapy. The most frequent malignancy was non-melanoma skin cancer (3.8%). Other second primary malignancies included solid tumors (including genitourinary and breast cancers) and melanoma. Advise patients to use sun protection and monitor for development of second primary malignancies.

Embryo-Fetal Toxicity: Based on animal findings, Jaypirca can cause fetal harm in pregnant women. Administration of pirtobrutinib to pregnant rats during organogenesis caused embryo-fetal toxicity, including embryo-fetal mortality and malformations at maternal exposures (AUC) approximately 3-times the recommended 200 mg/day dose. Advise pregnant women of potential risk to a fetus and females of reproductive potential to use effective contraception during treatment and for one week after last dose.

Adverse Reactions (ARs) in Patients with Mantle Cell Lymphoma Who Received Jaypirca
Serious ARs occurred in 38% of patients. Serious ARs occurring in ≥2% of patients were pneumonia (14%), COVID-19 (4.7%), musculoskeletal pain (3.9%), hemorrhage (2.3%), pleural effusion (2.3%), and sepsis (2.3%). **Fatal ARs** within 28 days of last dose of Jaypirca occurred in 7% of patients, most commonly due to infections (4.7%), including COVID-19 (3.1%).

Dose Modifications and Discontinuations: ARs led to dosage reductions in 4.7%, treatment interruption in 32%, and permanent discontinuation of Jaypirca in 9% of patients. ARs resulting in dosage modification in >5% of patients included pneumonia and neutropenia. ARs resulting in permanent discontinuation of Jaypirca in >1% of patients included pneumonia.

ARs (all Grades %; Grade 3-4 %) in ≥10% of Patients: fatigue (29; 1.6), musculoskeletal pain (27; 3.9), diarrhea (19; -), edema (18; 0.8), dyspnea (17; 2.3), pneumonia (16; 14), bruising (16; -), peripheral neuropathy (14; 0.8), cough (14; -), rash (14; -), fever (13; -), constipation (13; -), arthritis/arthralgia (12; 0.8), hemorrhage (11; 3.1), abdominal pain (11; 0.8), nausea (11; -), upper respiratory tract infections (10; 0.8), dizziness (10; -).

Select Laboratory Abnormalities (all Grades %; Grade 3 or 4 %) that Worsened from Baseline in ≥10% of Patients: hemoglobin decreased (42; 9), platelet count decreased (39; 14), neutrophil count decreased (36; 16), lymphocyte count decreased (32; 15), creatinine increased (30; 1.6), calcium decreased (19; 1.6), AST increased (17; 1.6), potassium decreased (13; 1.6), sodium decreased (13; -), lipase increased (12; 4.4), alkaline phosphatase increased (11; -), ALT increased (11; 1.6), potassium increased (11; 0.8). Grade 4 laboratory abnormalities in >5% of patients included neutrophils decreased (10), platelets decreased (7), lymphocytes decreased (6).

All grade ARs with higher frequencies in the total BRUIN population of patients with hematologic malignancies (n=583) were decreased neutrophil count (41%), bruising (20%), diarrhea (20%).

Drug Interactions
Strong CYP3A Inhibitors: Concomitant use with Jaypirca increased pirtobrutinib systemic exposure, which may increase risk of Jaypirca adverse reactions. Avoid use of strong CYP3A inhibitors during Jaypirca treatment. If concomitant use is unavoidable, reduce Jaypirca dosage according to the approved labeling.

Strong or Moderate CYP3A Inducers: Concomitant use with Jaypirca decreased pirtobrutinib systemic exposure, which may reduce Jaypirca efficacy. Avoid concomitant use of Jaypirca with strong or moderate CYP3A inducers. If concomitant use with moderate CYP3A inducers is unavoidable, increase the Jaypirca dosage according to the approved labeling.

Sensitive CYP2C8, CYP2C19, CYP3A, P-gP, BCRP Substrates: Concomitant use with Jaypirca increased their plasma concentrations, which may increase risk of adverse reactions related to these substrates for drugs that are sensitive to minimal concentration changes. Follow recommendations for these sensitive substrates in their approved labeling.

Use in Special Populations

Pregnancy and Lactation: Inform pregnant women of potential for Jaypirca to cause fetal harm. Verify pregnancy status in females of reproductive potential prior to starting Jaypirca and advise use of effective contraception during treatment and for one week after last dose. Presence of pirtobrutinib in human milk and effects on the breastfed child or on milk production is unknown. Advise women not to breastfeed while taking Jaypirca and for one week after last dose.

Geriatric Use: In the pooled safety population of patients with hematologic malignancies, 392 (67%) were ≥65 years of age. Patients aged ≥65 years experienced higher rates of Grade ≥3 ARs and serious ARs compared to patients <65 years of age.

Renal Impairment: Severe renal impairment (eGFR 15-29 mL/min) increases pirtobrutinib exposure. Reduce Jaypirca dosage in patients with severe

renal impairment according to the approved labeling. No dosage adjustment is recommended in patients with mild or moderate renal impairment.

PT HCP ISI MCL APP

Please see Prescribing Information and Patient Information for Jaypirca.

About Lilly
Lilly unites caring with discovery to create medicines that make life better for people around the world. We've been pioneering life-changing discoveries for nearly 150 years, and today our medicines help more than 47 million people across the globe. Harnessing the power of biotechnology, chemistry and genetic medicine, our scientists are urgently advancing new discoveries to solve some of the world's most significant health challenges, redefining diabetes care, treating obesity and curtailing its most devastating long-term effects, advancing the fight against Alzheimer's disease, providing solutions to some of the most debilitating immune system disorders, and transforming the most difficult-to-treat cancers into manageable diseases. With each step toward a healthier world, we're motivated by one thing: making life better for millions more people. That includes delivering innovative clinical trials that reflect the diversity of our world and working to ensure our medicines are accessible and affordable. To learn more, visit Lilly.com and Lilly.com/newsroom or follow us on Facebook, Instagram and LinkedIn. P-LLY

Jaypirca™ is a trademark owned by or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.

Verzenio® is a trademark owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.

Lilly Forward-Looking Statement
This press release contains forward-looking statements (as that term is defined in the Private Securities Litigation Reform Act of 1995) about Lilly's oncology portfolio and pipeline, including Verzenio, Jaypirca, LY3537982, and LY3410738, and reflects Lilly's current beliefs and expectations. However, as with any pharmaceutical product, there are substantial risks and uncertainties in the process of research, development, regulatory approval, and commercialization. Among other things, there can be no guarantee that future studies will be completed as planned, that future study results will be consistent with the results to date, that Verzenio, Jaypirca, LY3537982, and LY3410738 will receive (or receive additional) regulatory approvals or be commercially successful. For further discussion of these and other risks and uncertainties, see Lilly's most recent Form 10-K and Form 10-Q filings with the United States Securities and Exchange Commission. Except as required by law, Lilly undertakes no duty to update forward-looking statements to reflect events after the date of this release.

© Lilly USA, LLC 2023. ALL RIGHTS RESERVED.

Refer to: Lauren Cohen; lcohen@loxooncology.com; 617-678-2067 – media
Joe Fletcher; jfletcher@lilly.com; 317-296-2884 – investors

View original content to download multimedia:https://www.prnewswire.com/news-releases/lilly-announces-details-of-presentations-at-2023-american-association-for-cancer-research-aacr-annual-meeting-301771587.html

SOURCE Eli Lilly and Company