



Foundayo and Zepbound now covered for millions of Americans

May 28, 2026

CVS Caremark begins covering Foundayo June 1 and resumes Zepbound coverage by October 1

Eligible patients may pay as low as \$25 a month for their medicine

INDIANAPOLIS, May 28, 2026 /PRNewswire/ -- Eli Lilly and Company (NYSE: LLY) today announced an access milestone in obesity care: all three of the nation's largest pharmacy benefit managers (PBMs) will cover Lilly's full obesity medicine portfolio. This significantly expands the number of Americans who can access FDA-approved options through their existing insurance.

"For too long, effective obesity treatment has been out of reach for the people who need it," said Ilya Yuffa, executive vice president and president of Lilly USA and global customer capabilities. "Not all medicines work the same way for patients. Broader coverage puts real choice in the hands of millions of Americans and their doctors."

Lilly has worked across major PBMs to expand coverage for its obesity medicines. With today's news, coverage from CVS Caremark Commercial Template begins June 1 for Foundayo. For current Zepbound patients, coverage continues uninterrupted, and access will broaden across template plans by October 1. For eligible patients with commercial coverage, both medicines are available for as little as \$25 a month.* Medicare Part D beneficiaries may also be eligible to pay \$50 per month for their obesity medicines beginning July 1 through the Medicare GLP-1 Bridge program.**

Foundayo is the only once-daily GLP-1 pill for weight management that can be taken any time of day without food or water restrictions.¹ Zepbound is the most-prescribed injectable weight management medicine in the United States.² Together, they offer patients and their doctors treatment options that fit into people's lives.

To check coverage and explore savings options, visit [Foundayo.com](https://www.foundayo.com) or [Zepbound.com](https://www.zepbound.com). Coverage information is updated regularly, and people are encouraged to check back often to see the latest coverage and cost information. Talk to your healthcare provider about which treatment option may be right for you.

About Zepbound® (tirzepatide) injection

Zepbound® (tirzepatide) is the first and only dual GIP (glucose-dependent insulinotropic polypeptide) and GLP-1 (glucagon-like peptide-1) receptor agonist obesity medication. Zepbound tackles an underlying cause of excess weight. It reduces appetite and how much you eat. Zepbound is indicated for adults with obesity, or some adults who are overweight and also have at least one weight-related medical problem, to lose weight and keep it off. Additionally, Zepbound is FDA-approved to treat adults with moderate-to-severe obstructive sleep apnea and obesity. Zepbound should be used with a reduced calorie diet and increased physical activity.

About Foundayo™ (orforglipron)³

Foundayo™ (orforglipron) is FDA-approved for adults with obesity, or some adults with overweight who also have weight-related medical problems to reduce excess body weight and maintain weight reduction long term, alongside a reduced-calorie diet and increased physical activity. Foundayo is a once-daily small molecule (non-peptide) oral glucagon-like peptide-1 receptor agonist that can be taken any time of the day without restrictions on food and water intake.¹ Orforglipron was discovered by Chugai Pharmaceutical Co., Ltd. and licensed by Lilly in 2018. In addition to chronic weight management, orforglipron is being studied as a potential treatment for type 2 diabetes, obstructive sleep apnea, osteoarthritis knee pain, hypertension, peripheral artery disease and stress urinary incontinence.

Warnings - Foundayo and Zepbound may cause tumors in the thyroid, including thyroid cancer. Watch for possible symptoms, such as a lump or swelling in the neck, hoarseness, trouble swallowing, or shortness of breath. If you have any of these symptoms, tell your healthcare provider.

About ATTAİN-1 and ATTAİN-2 clinical trial program

The ATTAİN Phase 3 global clinical development program for orforglipron has enrolled more than 4,500 people with obesity or overweight across two global registration trials.

ATTAİN-1 (NCT05869903) is a Phase 3, 72-week, randomized, double-blind, placebo-controlled trial comparing the efficacy and safety of orforglipron at various doses to placebo in adults with obesity, or overweight with at least one of the following comorbidities: hypertension, dyslipidemia, obstructive sleep apnea or cardiovascular disease, who did not have diabetes. The trial randomized 3,127 participants across the U.S., Brazil, China, India, Japan, South Korea, Puerto Rico, Slovakia, Spain and Taiwan to receive various doses of orforglipron or placebo along with healthy diet and physical activity. The primary objective of the study was to demonstrate that orforglipron is superior to placebo in body weight reduction from baseline after 72 weeks.

ATTAİN-2 (NCT05872620) is a Phase 3, 72-week, randomized, double-blind, placebo-controlled trial comparing the efficacy and safety of various doses of orforglipron with placebo in adults with obesity or overweight and type 2 diabetes. The trial randomized over 1,600 participants across the U.S., Argentina, Australia, Brazil, China, Czechia, Germany, Greece, India, South Korea and Puerto Rico to receive various doses of orforglipron or placebo along with healthy diet and physical activity. The primary objective of the study was to demonstrate that orforglipron is superior to placebo in mean body weight change from baseline at 72 weeks.

Supported by the rigorous ATTAİN clinical trial program, Foundayo, along with diet and exercise, was proven to help people lose weight and keep it off. In the ATTAİN-1 trial, individuals taking the highest dose of Foundayo and who stayed on treatment lost an average of 27.3 pounds (12.4%) compared

to 2.2 pounds (0.9%) with placebo.⁴ Participants taking the highest dose of Foundayo, regardless of trial completion on study drug, lost an average of 25 pounds (11.1%), compared to 5.3 pounds (2.1%) with placebo.⁵

About SURMOUNT-1

Throughout the 72 week clinical trial, people who took Zepbound sustained weight loss—whether taking the 5 mg, 10 mg, or 15 mg dose along with diet and exercise. In a 72-week study of adults without diabetes, average weight loss was 15.0% (34 lbs) for 5 mg, 19.5% (44 lbs) for 10 mg, 20.9% (48 lbs) for 15 mg, and 3.1% (7 lbs) for placebo. Average starting weights were 226.8 lbs for 5 mg, 233.3 lbs for 10 mg, 232.8 lbs for 15 mg, and 231.0 lbs for placebo.

INDICATIONS AND SAFETY SUMMARY WITH WARNINGS

Zepbound® (ZEHP-bownd) is an injectable prescription medicine used with a reduced-calorie diet and increased physical activity to help adults with:

- obesity, or some adults with overweight who also have weight-related medical problems, to lose excess body weight and keep the weight off.
- moderate-to-severe obstructive sleep apnea (OSA) and obesity to improve their OSA.

Zepbound contains tirzepatide and should not be used with other tirzepatide-containing products or any GLP-1 receptor agonist medicines. It is not known if Zepbound is safe and effective for use in children.

Warnings - Zepbound may cause tumors in the thyroid, including thyroid cancer. Watch for possible symptoms, such as a lump or swelling in the neck, hoarseness, trouble swallowing, or shortness of breath. If you have any of these symptoms, tell your healthcare provider.

- Do not use Zepbound if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC).
- Do not use Zepbound if you have Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).
- Do not use Zepbound if you have had a serious allergic reaction to tirzepatide or any of the ingredients in Zepbound.

KwikPen®: Do not share your KwikPen with other people, even if the pen needle has been changed. You may give other people a serious infection or get a serious infection from them.

Zepbound may cause serious side effects, including:

Severe stomach problems. Stomach problems, sometimes severe, have been reported in people who use Zepbound. Tell your healthcare provider if you have stomach problems that are severe or will not go away.

Dehydration leading to kidney problems. Diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration), which may cause kidney problems. It is important for you to drink fluids to help reduce your chance of dehydration. Tell your healthcare provider right away if you have nausea, vomiting, or diarrhea that does not go away.

Gallbladder problems. Gallbladder problems have happened in some people who use Zepbound. Tell your healthcare provider right away if you get symptoms of gallbladder problems, which may include pain in your upper stomach (abdomen), fever, yellowing of skin or eyes (jaundice), or clay-colored stools.

Inflammation of the pancreas (pancreatitis). Stop using Zepbound and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back.

Serious allergic reactions. Stop using Zepbound and get medical help right away if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue or throat, problems breathing or swallowing, severe rash or itching, fainting or feeling dizzy, or very rapid heartbeat.

Low blood sugar (hypoglycemia). Your risk for getting low blood sugar may be higher if you use Zepbound with medicines that can cause low blood sugar, such as a sulfonylurea or insulin. **Signs and symptoms of low blood sugar** may include dizziness or light-headedness, sweating, confusion or drowsiness, headache, blurred vision, slurred speech, shakiness, fast heartbeat, anxiety, irritability, mood changes, hunger, weakness or feeling jittery.

Changes in vision in patients with type 2 diabetes. Tell your healthcare provider if you have changes in vision during treatment with Zepbound.

Food or liquid getting into the lungs during surgery or other procedures that use anesthesia or deep sleepiness (deep sedation). Zepbound may increase the chance of food getting into your lungs during surgery or other procedures. Tell all your healthcare providers that you are taking Zepbound before you are scheduled to have surgery or other procedures.

Common side effects

The most common side effects of Zepbound include nausea, diarrhea, vomiting, constipation, stomach (abdominal) pain, indigestion, injection site reactions, feeling tired, allergic reactions, belching, hair loss, and heartburn. These are not all the possible side effects of Zepbound. Talk to your healthcare provider about any side effect that bothers you or doesn't go away.

Tell your doctor if you have any side effects. **You can report side effects at 1-800-FDA-1088 or www.fda.gov/medwatch.**

Before using Zepbound

- **Your healthcare provider should show you how to use Zepbound before you use it for the first time.**
- **Talk to your healthcare provider about low blood sugar and how to manage it. Tell your healthcare provider if you are taking medicines to treat diabetes including an insulin or sulfonylurea.**
- **If you take birth control pills by mouth, talk to your healthcare provider before you use Zepbound. Birth control**

pills may not work as well while using Zepbound. Your healthcare provider may recommend another type of birth control for 4 weeks after you start Zepbound and for 4 weeks after each increase in your dose of Zepbound.

Review these questions with your healthcare provider:

- Do you have other medical conditions, including problems with your pancreas, or severe problems with your stomach, such as slowed emptying of your stomach (gastroparesis) or problems digesting food?
- Do you take diabetes medicines, such as insulin or sulfonylureas?
- Do you have a history of diabetic retinopathy?
- Are you scheduled to have surgery or other procedures that use anesthesia or deep sleepiness (deep sedation)?
- Do you take any other prescription medicines or over-the-counter drugs, vitamins, or herbal supplements?
- Are you pregnant, plan to become pregnant, breastfeeding, or plan to breastfeed? Zepbound may harm your unborn baby. Tell your healthcare provider if you become pregnant while using Zepbound. Zepbound may pass into your breast milk. You should talk with your healthcare provider about the best way to feed your baby while using Zepbound.

- **Pregnancy Exposure Registry:** There will be a pregnancy exposure registry for women who have taken Zepbound during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry, or you may contact Lilly at 1-800-LillyRx (1-800-545-5979).

How to take

- Read the Instructions for Use that come with Zepbound.
- Use Zepbound exactly as your healthcare provider says.
- Use Zepbound with a reduced-calorie diet and increased physical activity.
- Inject Zepbound under the skin (subcutaneously) of your stomach (abdomen), thigh, or have another person inject in the back of the upper arm. **Do not** inject ZEPBOUND into a muscle (intramuscularly) or vein (intravenously).
- **Use Zepbound 1 time each week, at any time of the day.**
- Change (rotate) your injection site with each weekly injection. **Do not** use the same site for each injection.
- If you take too much Zepbound, call your healthcare provider, call the Poison Help line at 1-800-222-1222 or go to the nearest hospital emergency room right away.

If you take too much Zepbound, call your healthcare provider, call the Poison Help line at 1-800-222-1222 or go to the nearest hospital emergency room right away.

Zepbound is approved as a 2.5 mg, 5 mg, 7.5 mg, 10 mg, 12.5 mg, and 15 mg injection.

Learn more

Zepbound is a prescription medicine. For more information, call 1-800-LillyRx (1-800-545-5979) or go to www.zepbound.lilly.com.

This summary provides basic information about Zepbound but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your healthcare provider. Be sure to talk to your healthcare provider about Zepbound and how to take it. Your healthcare provider is the best person to help you decide if Zepbound is right for you.

ZP CON BS 25FEB2026

Zepbound[®], its delivery device base and KwikPen[®] are registered trademarks owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.

INDICATION AND SAFETY SUMMARY WITH WARNINGS

Foundayo[™] (fown-DAY-oh) is a prescription medicine used with a reduced-calorie diet and increased physical activity to help adults with obesity, or some adults with overweight who also have weight-related medical problems, to lose excess body weight and keep the weight off.

- Foundayo should not be used with other GLP-1 receptor agonist medicines.
- It is not known if Foundayo is safe and effective for use in children.

Warnings – Foundayo may cause tumors in the thyroid, including thyroid cancer. Watch for possible symptoms, such as a lump or swelling in the neck, hoarseness, trouble swallowing, or shortness of breath. If you have any of these symptoms, tell your healthcare provider.

- Do not use Foundayo if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC).
- Do not use Foundayo if you have Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).
- Do not use Foundayo if you have had a serious allergic reaction to orforglipron or any of the ingredients in Foundayo.

Foundayo may cause serious side effects, including:

Inflammation of the pancreas (pancreatitis). Stop taking Foundayo and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without nausea or vomiting. Sometimes you may feel the pain from your abdomen to your back.

Severe stomach problems. Stomach problems, sometimes severe, have been reported in people who use Foundayo. Tell your healthcare provider if you have stomach problems that are severe or will not go away.

Dehydration leading to kidney problems. Diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration), which may cause kidney problems. It is important for you to drink fluids to help reduce your chance of dehydration. Tell your healthcare provider right away if you have nausea, vomiting, or diarrhea that does not go away.

Low blood sugar (hypoglycemia). Your risk for getting low blood sugar may be higher if you use Foundayo with medicines that can cause low blood sugar, such as an insulin or sulfonylurea. **Signs and symptoms of low blood sugar may include** dizziness or light-headedness, sweating, confusion or drowsiness, headache, blurred vision, slurred speech, shakiness, fast heartbeat, anxiety, irritability, mood changes, hunger, weakness, or feeling jittery.

Serious allergic reactions. Stop using Foundayo and get medical help right away if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue or throat, problems breathing or swallowing, severe rash or itching, fainting or feeling dizzy, or very rapid heartbeat.

Changes in vision in patients with type 2 diabetes. Tell your healthcare provider if you have changes in vision during treatment with Foundayo.

Gallbladder problems. Gallbladder problems have happened in some people who use Foundayo. Tell your healthcare provider right away if you get symptoms of gallbladder problems, which may include pain in your upper stomach (abdomen), fever, yellowing of skin or eyes (jaundice), or clay-colored stools.

Food or liquid getting into the lungs during surgery or other procedures that use anesthesia or deep sleepiness (deep sedation). Foundayo may increase the chance of food getting into your lungs during surgery or other procedures. Tell your healthcare providers that you are taking Foundayo before you are scheduled to have surgery or other procedures.

Common side effects

The most common side effects of Foundayo include nausea, constipation, diarrhea, vomiting, indigestion, stomach (abdominal) pain, headache, swollen belly, feeling tired, belching, heartburn, gas, and hair loss. These are not all the possible side effects of Foundayo. Talk to your healthcare provider about any side effect that bothers you or doesn't go away.

Tell your doctor if you have any side effects. **You can report side effects at 1-800-FDA-1088 or www.fda.gov/medwatch.**

Before taking Foundayo

- **Tell your healthcare provider about all the medicines you take.** Foundayo may affect the way some medicines work, and some medicines may affect the way Foundayo works.
- **Pregnancy Exposure Registry:** There will be a pregnancy exposure registry for women who have taken Foundayo during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry, or you may contact Eli Lilly and Company at 1-800-LillyRx (1-800-545-5979).
- **If you take birth control pills by mouth, talk to your healthcare provider before you take Foundayo. Birth control pills may not work as well while taking Foundayo.** Your healthcare provider may recommend another type of birth control for 30 days after starting Foundayo and for 30 days after each dose increase of Foundayo.
- **Talk to your healthcare provider about low blood sugar and how to manage it.** Tell your healthcare provider if you are taking medicines to treat diabetes including an insulin or sulfonylurea.

Review these questions with your healthcare provider:

- Do you have other medical conditions, including problems with your pancreas or kidneys, or severe problems with your liver, severe problems with your stomach, such as slowed emptying of your stomach (gastroparesis) or problems digesting food?
- Do you have a history of diabetic retinopathy?
- Are you scheduled to have surgery or other procedures that use anesthesia or deep sleepiness (deep sedation)?
- Are you pregnant or plan to become pregnant? Foundayo may harm your unborn baby.
- Are you breastfeeding or plan to breastfeed? Breastfeeding is not recommended during treatment with Foundayo.
- Do you take any other prescriptions or over-the-counter medicines, vitamins, or herbal supplements?

How to take

- Take Foundayo exactly as your healthcare provider tells you to.
- Use Foundayo with a reduced-calorie diet and increased physical activity.
- Take Foundayo by mouth 1 time each day, with or without food.
- Swallow tablets whole. Do not break, crush, or chew the tablet.
- If you miss a dose, take it as soon as possible. **Do not take 2 doses of Foundayo in the same day.**
- **Do not take more than 1 tablet per day.**
- If you miss taking Foundayo for 7 or more days in a row, call your healthcare provider to talk about how to restart your treatment.
- If you take too much Foundayo, call your healthcare provider or Poison Help line at 1-800-222-1222 or go to the nearest hospital emergency room right away.

Learn more

Foundayo is a prescription medicine available in 0.8 mg, 2.5 mg, 5.5 mg, 9 mg, 14.5 mg, or 17.2 mg oral tablets. For more information, call 1-800-545-5979 or go to foundayo.lilly.com.

This summary provides basic information about Foundayo but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your doctor. Be sure to talk to your doctor or other healthcare provider about Foundayo and how to take it. Your doctor is the best person to help you decide if Foundayo is right for you.

OG CON BS APR2026

Foundayo™ is a trademark of Eli Lilly and Company.

Endnotes and References

*Terms and conditions apply. For Zepbound, available with auto-injector presentation.

**For further details regarding Medicare GLP-1 Bridge program eligibility, visit [CMS.gov](https://www.cms.gov).

1. Ma X, Liu R, Pratt EJ, Benson CT, Bhattachar SN, Sloop KW. Effect of Food Consumption on the Pharmacokinetics, Safety, and Tolerability of Once-Daily Orally Administered Orforglipron (LY3502970), a Non-peptide GLP-1 Receptor Agonist. *Diabetes Ther.* 2024 Apr;15(4):819-832. <https://doi.org/10.1007/s13300-024-01554-1>. Epub 2024 Feb 24. PMID: 38402332; PMCID: PMC10951152.
2. Based on IQVIA® National Prescription Audit Data, representing 94% of US prescription data as of January 10, 2025.
3. Foundayo. Prescribing Information. Lilly USA, LLC.
4. The efficacy estimand represents efficacy had all randomized participants remained on study intervention (with possible dose interruptions and modifications) for 72 weeks without initiating prohibited weight management treatments.³
5. The treatment-regimen estimand represents the estimated average treatment effect regardless of adherence to study intervention or initiation of prohibited weight management treatments.³

About Lilly

Lilly is a medicine company turning science into healing to make life better for people around the world. We've been pioneering life-changing discoveries for 150 years, and today our medicines help tens of millions of people across the globe. Harnessing the power of biotechnology, chemistry and genetic medicine, our scientists are urgently advancing new discoveries to solve some of the world's most significant health challenges: redefining diabetes care; treating obesity and curtailing its most devastating long-term effects; advancing the fight against Alzheimer's disease; providing solutions to some of the most debilitating immune system disorders; and transforming the most difficult-to-treat cancers into manageable diseases. With each step toward a healthier world, we're motivated by one thing: making life better for millions more people. That includes delivering innovative clinical trials that reflect the diversity of our world and working to ensure our medicines are accessible and affordable. To learn more, visit [Lilly.com](https://www.lilly.com) and [Lilly.com/news](https://www.lilly.com/news), or follow us on [Facebook](https://www.facebook.com/lilly), [Instagram](https://www.instagram.com/lilly), and [LinkedIn](https://www.linkedin.com/company/lilly). P-LLY

Trademarks and Trade Names

All trademarks or trade names referred to in this press release are the property of the company, or, to the extent trademarks or trade names belonging to other companies are referenced in this press release, the property of their respective owners. Solely for convenience, the trademarks and trade names in this press release are referred to without the ® and ™ symbols, but such references should not be construed as any indicator that the company or, to the extent applicable, their respective owners will not assert, to the fullest extent under applicable law, the company's or their rights thereto. We do not intend the use or display of other companies' trademarks and trade names to imply a relationship with, or endorsement or sponsorship of us by, any other companies.

Cautionary Statement Regarding Forward-Looking Statements

This press release contains forward-looking statements (as that term is defined in the Private Securities Litigation Reform Act of 1995), including statements about the supply and access of Zepbound (tirzepatide) as a treatment for adults with obesity or overweight and Foundayo as a treatment for adults with obesity or some adults with overweight who also have weight-related medical problems and reflects Lilly's current belief and expectations. However, as with any pharmaceutical product, there are substantial risks and uncertainties in the process of drug research, development, access, and commercialization. Among other things, there can be no guarantee that future study results will be consistent with the results to date, that Zepbound will receive additional regulatory approvals, or that Lilly will execute its access and other strategies as planned. For further discussion of these and other risks and uncertainties, see Lilly's most recent Form 10-K and Form 10-Q filings with the United States Securities and Exchange Commission. Except as required by law, Lilly undertakes no duty to update forward-looking statements to reflect events after the date of this release.

CMAT-33004 5/2026 ©Lilly USA, LLC 2026. All rights reserved.

Refer to: Rachel Sorvig; sorvig_rachel@lilly.com ; 317-607-7507 (Media)

Michael Czapar; czapar_michael_c@lilly.com; 317-617-0983 (Investors)

Lilly

View original content to download multimedia: <https://www.prnewswire.com/news-releases/foundayo-and-zepbound-now-covered-for-millions-of-americans-302784360.html>

SOURCE Eli Lilly and Company